

Some Loci in the Animation of a Sangaku Diagram

Junghyun Lee, Minyoung Hwang, and Cheolwon Bae

Abstract. In a symmetric partition of a regular n -gon into n congruent subtriangles and a regular n -gon in the center, we determine the loci of the incenter and points of tangency of the incircle a subtriangle.

A famous Sangaku problem ([1, Problem 2.1.7], [2]) asks to partition an equilateral triangle into four subtriangles with congruent incircle (see Figure 1). Ito and Wimmer [3] considered the same problem for general regular polygons (see Figure 2 for the case of a regular pentagon). In this note, we consider a dynamic situation by letting the congruent subtriangles by the sides of the regular polygon vary, and examine the loci of various points in the configuration.




Figure 1




Figure 2

Given a regular n -gon $\mathcal{P} := A_1A_2 \cdots A_n$, $k = 1, 2, \dots, n$, with center O , pass a line $\ell_k(\theta)$ through the vertex A_k such that the directed angle $(A_kA_{k+1}, \ell_k(\theta)) = \theta \in (0, \pi - \frac{2\pi}{n})$. Here indices are taken modulo n so that $A_{n+1} = A_1$ etc. Let $A'_k(\theta)$ be the intersection of $\ell_k(\theta)$ and $\ell_{k+1}(\theta)$ (see Figure 3). When there is no danger of confusion, we shall simply write A'_k for $A'_k(\theta)$. Then the regular n -gon \mathcal{P} is partitioned into

- (i) n congruent triangles $A'_kA_kA_{k+1}$ for $k = 1, 2, \dots, n$, and
- (ii) a regular n -gon $\mathcal{P}' = A'_1A'_2 \cdots A'_n$ at the center.




Figure 3

It is clear that the center of the regular n -gon $A'_1 A'_2 \cdots A'_n$ is the fixed point O . On the other hand, the locus of $A'_k(\theta)$ is the part of the circle $OA_k A_{k+1}$ in the interior of \mathcal{P} . This is because

$$\angle A_k A'_k A_{k+1} = \left(\frac{2\pi}{n} + \theta \right) - \theta = \frac{2\pi}{n} = \angle A_k O A_{k+1},$$

from which $A_k, A'_k, O,$ and A_{k+1} are concyclic.

Suppose the incircle of \mathcal{P}' touches the sides $A'_{k-1} A'_k$ at B_k , the midpoint of A'_{k-1} and A'_k on l_k . Since OB_k is perpendicular to $A'_{k-1} A'_k$, it is perpendicular the line $l_k(\theta)$ through A_k . Therefore, $OB_k A_k$ is a right angle, and B_k lies on the part of the circle with diameter OA_k inside the regular polygon \mathcal{P} . If M_{k-1} and M_k are the midpoints of $A_{k-1} A_k$ and $A_k A_{k+1}$, then this is the arc of the circle $M_{k-1} O M_k$ in the interior of \mathcal{P} .

Now we consider the incircle of triangle $\mathbf{T}_k = A'_k A_k A_{k+1}$, with incenter I_k , and tangent to l_k, l_{k+1} at the points $T_{k,k}, T_{k+1,k}$ respectively.

Note that

$$\angle A_k I_k A_{k+1} = \frac{\theta}{2} + \frac{2\pi}{n} + \frac{1}{2} \left(\pi - \frac{2\pi}{n} - \theta \right) = \frac{\pi}{2} - \frac{\pi}{n}$$

is independent of θ . This means that the locus of I_k is the part of a circle through A_k and A_{k+1} in the interior of \mathcal{P} . Its center C_k is the intersection of the perpendicular bisector of $A_k A_{k+1}$ and the *external* bisector of angle $A_{k-1} A_k A_{k+1}$.

We summarize these simple results in the following proposition.

Proposition 1. Let $\mathcal{P} := A_1A_2 \cdots A_n$ be a regular n -gon with center O . For $k = 1, 2, \dots, n$, let $\ell_k(\theta)$ be the line through the vertex A_k such that the directed angle $(A_kA_{k+1}, \ell_k(\theta)) = \theta$ (with indices taken modulo n). As θ varies in $(0, \pi - \frac{2\pi}{n})$, the loci of

- (a) the intersection $A'_k(\theta)$ of $\ell_k(\theta)$ and $\ell_{k+1}(\theta)$ is the part of the circle OA_kA_{k+1} in the interior of \mathcal{P} ,
- (b) the point of tangency B_k of the incircle of the regular n -gon $A'_1A'_2 \cdots A'_n$ with the line $\ell_k(\theta)$ is the part of the circle $OM_{k-1}M_k$ in the interior of \mathcal{P} , M_k being the midpoint A_kA_{k+1} ,
- (c) the incenter I_k of triangle $A'_kA_kA_{k+1}$ is the arc of the circle, center C_k , passing through A_{k-1} , C_k being the intersection of the perpendicular bisector of A_kA_{k+1} and the external bisector of angle $A_{k-1}A_kA_{k+1}$.

We compute some of the lengths in this configuration. In Figure 3, let a be the length of a side of the regular n -gon \mathcal{P} . Suppose in triangle $A'_kA_kA_{k+1}$, $A'_kA_{k+1} = b$ and $A_kA'_k = c$. By the law of sines,

$$b = \frac{a}{\sin \frac{2\pi}{n}} \sin \theta,$$

$$c = \frac{a}{\sin \frac{2\pi}{n}} \sin \left(\frac{2\pi}{n} + \theta \right).$$

Theorem 2. For $k = 1, 2, \dots, n$, let $T_{k,k}$ and $T_{k+1,k}$ be the points of tangency of the incircle of triangle $A'_kA_kA_{k+1}$ with the lines $\ell_k(\theta)$ and $\ell_{k+1}(\theta)$ respectively. θ varies in $(0, \pi - \frac{2\pi}{n})$, the loci of $T_{k,k}$ and $T_{k+1,k}$ are the parts of limaçon inside the regular n -gon \mathcal{P} , symmetric with respect to the perpendicular bisector of A_kA_{k+1} .

Proof. The point of tangency $T_{k,k}$ is the point on $\ell_k(\theta)$ uniquely determined by the length of $A_kT_{k,k}$. Now, in triangle $A'_kA_kA_{k+1}$,

$$\begin{aligned} A_kT_{k,k} &= \frac{1}{2}(a + c - b) \\ &= \frac{a}{2} + \frac{a}{2 \sin \frac{2\pi}{n}} \left(\sin \left(\frac{2\pi}{n} + \theta \right) - \sin \theta \right) \\ &= \frac{a}{2} + \frac{a}{2 \sin \frac{2\pi}{n}} \cdot 2 \cos \left(\frac{\pi}{n} + \theta \right) \sin \frac{\pi}{n} \\ &= \frac{a}{2} + \frac{a}{2 \cos \frac{\pi}{n}} \cdot \cos \left(\frac{\pi}{n} + \theta \right). \end{aligned}$$

Let C_k be the intersection of the perpendicular bisector of A_kA_{k+1} and the external angle of $A_{k-1}A_kA_{k+1}$. Note that $A_kC_k = \frac{a}{2 \cos \frac{\pi}{n}}$. In a polar coordinate system with pole at A_k and polar axis the half line A_kA_{k+1} , the equation

$$\rho = \frac{a}{2 \cos \frac{\pi}{n}} \cos \left(\frac{\pi}{n} + \theta \right)$$

represents the circle with A_kC_k as diameter. Therefore,

$$\rho = \frac{a}{2} + \frac{a}{2 \cos \frac{\pi}{n}} \cos \left(\frac{\pi}{n} + \theta \right)$$

is a limaçon: If the circle with diameter $A_k C_k$ intersects $\ell_k(\theta)$ at D_k , then $T_{k,k}$ is the point obtained by translating D_k by $\frac{a}{2}$ (along $A_k D_k$); see Figure 4 for the case of a regular pentagon $A_1 A_2 A_3 A_4 A_5$ with $k = 1$.




Figure 4

The locus of $T_{k+1,k}$ is clearly the reflection of the locus of $T_{k,k}$ in the perpendicular bisector of $A_k A_{k+1}$. It is determined by

$$A_{k+1} T_{k+1,k} = \frac{1}{2}(a - c + b) = \frac{a}{2} - \frac{a}{2 \cos \frac{\pi}{n}} \cdot \cos \left(\frac{\pi}{n} + \theta \right).$$

This is the limaçon with respect to the circle diameter $A_{k+1} C_k$ and length $\frac{a}{2}$. \square

References

[1] H. Fukagawa and D. Pedoe, *Japanese Temple Geometry Problems*, Charles Babbage Research Centre, Winnipeg, 1989.
 [2] H. Fukagawa and T. Rothman, *Sacred Mathematics*, Princeton University Press, 2008.
 [3] N. Ito and H. Wimmer, H, A Sangaku-type problem with regular polygons, triangles, and congruent incircles, *Forum Geom.*, 13 (2013) 185–190.

Junghyun Lee: Changwon Science High School, 159 beon-gil 30, Pyeongsan-ro, Uichang-gu,
Changwon-si, Gyeongsangnam-do, Korea
E-mail address: science0229@gmail.com

Minyoung Hwang: Changwon Science High School, 159 beon-gil 30, Pyeongsan-ro, Uichang-gu,
Changwon-si, Gyeongsangnam-do, Korea
E-mail address: myhwang99@naver.com

Cheolwon Bae: Changwon Science High School, 159 beon-gil 30, Pyeongsan-ro, Uichang-gu,
Changwon-si, Gyeongsangnam-do, Korea
E-mail address: cw2184@naver.com